

On the modularity of Calabi-Yau threefolds containing elliptic ruled surfaces

K. Hulek and H. Verrill

Abstract

In this paper we prove that (not necessarily rigid) Calabi-Yau threefolds defined over $\mathbb{Q}$ which contain sufficiently many elliptic ruled surfaces are modular (under certain mild restrictions on the primes of bad reduction). This is an application of the results of Dieulefait and Manoharmayum [DM], [D] who showed modularity for rigid Calabi-Yau threefolds.

0 Introduction

In the last two years different authors found several examples of non-rigid Calabi-Yau varieties defined over $\mathbb{Q}$ for which modularity could be proved (see [LY], [HV] and [Schu2]). The examples of Livné and Yui arise by taking a quotient of a product of a $K3$ surface with an elliptic curve, whereas the examples constructed by Schütt and the authors have the structure of a fibre product of elliptic surfaces. In each case the motive splits into low dimensional pieces which can be explained geometrically.

In this brief note we want to give a criterion which implies modularity if the Calabi-Yau variety contains sufficiently many elliptic ruled surfaces. This applies to the examples given in [HV] and [Schu2] and can also be used to explain the part of the motive in the examples in [LY] which does not come from the transcendental lattice of the $K3$ surface. The main ingredient of this paper is an application of a modularity result proved by Dieulefait and Manoharmayum in their paper [DM] (and its extension by Dieulefait [D]) where they establish modularity criteria for rigid Calabi-Yau threefolds.

Finally we shall show how to use our criterion to construct numerous examples of modular (birational) Calabi-Yau varieties. In some of these examples we have strong numerical evidence as to the modular forms which appear. The levels include 10,12,16,17,21,27,28,32,35,48,55,60,64,68,90,480 and the corresponding varieties are listed in Tables 1,2, 3 and 4.

Acknowledgements. We are very grateful to Professors Dieulefait and Manoharmayum for the exchange of some e-mails in which they explained some of their results and details about l -adic Galois representations to us.

We would like to thank the referee for computing the levels of the weight 4 modular forms appearing in Tables 2, and 3, and the level 480 example in Table 1. We also thank Matthias Schuett for reading through our manuscript and making several suggestions for improvements as well as for computational help with one example.

1 A modularity criterion

Let X be a smooth projective variety with $h^{30}(X) = h^{03}(X) = 1$. (In our applications this will be a Calabi-Yau variety or a variety birational to a nodal Calabi-Yau variety.) A (birational) elliptic ruled surface Y is a smooth projective surface which is birational to a $\mathbb{P}^1$ -bundle over an elliptic curve. In this case $h^{21}(Y) = h^{12}(Y) = 1$ and $h^{30}(Y) = h^{03}(Y) = 0$.

Definition 1.1. Let X be smooth projective and define $b := h^{12}(X)$. Let $Y_j, j = 1, \dots, b$ be birational elliptic ruled surfaces contained in X . We say that these surfaces *span* $H^{21}(X) \oplus H^{12}(X)$ if the natural map

$$H^3(X, \mathbb{C}) \rightarrow \oplus_{j=1}^b H^3(Y_j, \mathbb{C})$$

is surjective.

Remark 1.2. If the surfaces Y_j are as in the previous definition, then we have an exact sequence

$$0 \rightarrow H^{30}(X) \oplus H^{03}(X) \rightarrow H^3(X, \mathbb{C}) \rightarrow \oplus_{j=1}^b H^3(Y_j, \mathbb{C}) \rightarrow 0.$$

This follows since the restriction of a $(3, 0)$ or a $(0, 3)$ form to an elliptic ruled surface must vanish and together with the assumption that the surfaces Y_j span $H^{21}(X) \oplus H^{12}(X)$ this gives the claim. In particular we obtain an isomorphism $H^{21}(X) \oplus H^{12}(X) \cong \oplus_{j=1}^b H^3(Y_j, \mathbb{C}) = \oplus_{j=1}^b (H^{21}(Y_j) \oplus H^{12}(Y_j))$.

Remark 1.3. If X is a smooth Calabi-Yau threefold and Y is a smooth elliptic ruled surface contained in X , then a straightforward argument from intersection theory (cf. the proof of Proposition 2.1) using the adjunction formula shows that the map $H^3(X, \mathbb{C}) \rightarrow H^3(Y, \mathbb{C})$ is surjective. If $h^{12}(X) = 1$ then this shows that we are automatically in the situation described above. In fact this observation is often useful when one wants to prove modularity of Calabi-Yau threefolds with $h^{12}(X) = 1$.

We now assume that we are in an arithmetic situation, i.e. that the variety X as well as the subvarieties Y_j are defined over $\mathbb{Q}$. When we speak about a birational elliptic surface Y defined over $\mathbb{Q}$ we mean that there is a birational map defined over $\mathbb{Q}$ from Y to $E \times \mathbb{P}^1$. In what follows the assumption that all Y_j are defined over $\mathbb{Q}$ can often be weakened to

the assumption that the union of the surfaces Y_j is defined over $\mathbb{Q}$, but for the sake of simplicity we shall assume that all the Y_j themselves are defined over $\mathbb{Q}$. Then the map $H^3(X) \rightarrow \bigoplus_{j=1}^b H^3(Y_j)$ can be interpreted in the l -adic cohomology and gives rise to a morphism of $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ -modules $H_{\text{ét}}^3(X, \mathbb{Q}_l) \rightarrow \bigoplus H_{\text{ét}}^3(Y_j, \mathbb{Q}_l)$. (Strictly speaking we have a representation of the Galois group $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ on the dual vector spaces, but we shall, by misuse of language, refer to the cohomology groups as Galois representations.) If the surfaces Y_j span $H^{21}(X) \oplus H^{12}(X)$ this gives rise to an exact sequence of $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ -modules

$$0 \rightarrow U \rightarrow H_{\text{ét}}^3(X, \mathbb{Q}_l) \rightarrow \bigoplus H_{\text{ét}}^3(Y_j, \mathbb{Q}_l) \rightarrow 0 \quad (1)$$

where U has dimension 2.

We shall want to make use of a criterion proved by Dieulefait and Manoharmayum [DM] and further extended by Dieulefait [D] which was developed in order to prove the modularity of rigid Calabi-Yau varieties.

Proposition 1.4. Let l be a prime and let

$$\rho : \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}) \rightarrow \text{GL}_2(\overline{\mathbb{Q}}_l)$$

be an irreducible continuous Galois representation, unramified outside finitely many primes, and having determinant ε_l^3 where ε_l is the cyclotomic character. Furthermore assume the following:

- (1) ρ is crystalline at l with Hodge-Tate weights $\{0, 3\}$.
- (2) The residual representation

$$\overline{\rho} : \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}) \rightarrow \text{GL}_2(\overline{\mathbb{F}}_l)$$

takes values in $\text{GL}_2(\mathbb{F}_l)$.

Moreover assume that one of the following assumptions holds:

- (DM1) $l = 7$ and ρ is unramified at 3.
- (DM2) $l = 5$ and there is a prime $p \equiv \pm 2 \pmod{5}$ where ρ is unramified and the trace of Frobenius $t_3(p)$ at p is not divisible by 5, where we define $t_3(p) := \text{trace}(\rho(\text{Frob}_p))$.
- (D) $l > 3$, ρ is unramified at 3 and $t_3(3)$ is not divisible by 3.

Then ρ is modular.

Proof. Dieulefait and Manoharmayum proved that either of the conditions (DM1) or (DM2) are sufficient for modularity (see [DM, Remark 3.2]). The fact that condition (D) is sufficient was shown by Dieulefait ([D, Theorem 1.1]). $\square$

We now want to apply this to our situation.

Proposition 1.5. Let X be a smooth projective threefold defined over $\mathbb{Q}$ with $h^{30}(X) = h^{03}(X) = 1$ and assume that X contains birational ruled surfaces $Y_j, j = 1, \dots, b$ which are defined over $\mathbb{Q}$ and which span $H^{21}(X) \oplus H^{12}(X)$. Let ρ be the 2-dimensional Galois representation given by the kernel U from the exact sequence

$$0 \rightarrow U \rightarrow H_{\text{ét}}^3(X, \mathbb{Q}_l) \rightarrow \oplus H_{\text{ét}}^3(Y_j, \mathbb{Q}_l) \rightarrow 0.$$

Assume that one of the following conditions holds:

- (i) X has good reduction at 3 and 7 or
- (ii) X has good reduction at 5 and some at prime $p \equiv \pm 2 \pmod{5}$ and the trace of Frob_p on U is not divisible by 5 or
- (iii) X has good reduction at 3 and the trace of Frob_3 on U is not divisible by 3.

Then X is modular. More precisely

$$L(X, s) \stackrel{\circ}{=} L(f_4, s) \prod_{j=1}^b L(g_2^j, s-1) \quad (2)$$

where f_4 is a weight 4 form and the g_2^j are the weight 2 forms associated to the base curves E_j of the birational ruled surfaces Y_j . (As usual $\stackrel{\circ}{=}$ denotes equality of the L -series upto possibly finitely many factors.)

Proof. Since the surfaces Y_j are birational over $\mathbb{Q}$ to a product $E_j \times \mathbb{P}^1$ we have

$$L(Y_j, s) \stackrel{\circ}{=} L(g_2^j, s-1)$$

where g_2^j is the weight 2 form associated to the elliptic curve E_j . After passing to the semi-simplification we have an isomorphism

$$H_{\text{ét}}^3(X, \mathbb{Q}_l) \cong U \oplus \bigoplus_{j=1}^b H_{\text{ét}}^3(Y_j, \mathbb{Q}_l).$$

Our claim follows if we can show that we can apply the results of Dieulefait and Manoharmayum, resp. Dieulefait to deduce the modularity of the 2-dimensional $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ representation U . Clearly U is a continuous representation which is unramified outside finitely many primes, namely the primes of bad reduction. It follows from the Weil conjectures as proved by Deligne that U is irreducible [DM, pp. 1,2]. Moreover if U is unramified at l then its determinant is ε_l^3 . Condition (2) of 1.4 follows since U is a subspace

of $H_{\text{ét}}^3(X, \mathbb{Q}_l)$ for which (2) holds. Similarly we can deduce that U is crystalline and has Hodge-Tate weights $\{0, 3\}$. The first follows e.g. from the fact that $H_{\text{ét}}^3(X, \mathbb{Q}_l)$ is crystalline for $l > 3$ (see [FM, Theorem 2.2 A (ii)]) and the fact that this property is preserved for submodules and quotients (see [FM, Proposition 1.7]). The statement about the Hodge-Tate weight follows from standard comparison theorems (see [FM, Section 3.1]). The conditions (DM1), (DM2) or (D) follow immediately from the hypothesis of the theorem. $\square$

Remark 1.6. The weight 4 form which is associated to the 2-dimensional representation U can in many cases be determined explicitly by the method of Faltings, Livné and Serre.

2 Fibre products

We shall now investigate a certain class of Calabi-Yau threefolds which often (but not always) contain many elliptic ruled surfaces in the sense of our discussion in the previous section.

Let $r : Y \rightarrow \mathbb{P}^1$ and $r' : Y' \rightarrow \mathbb{P}^1$ be semi-stable rational elliptic fibrations with a section. Then the fibre product

$$W = Y \times_{\mathbb{P}^1} Y'$$

is a nodal Calabi-Yau variety. We denote by $\hat{W}$ a small and by $\tilde{W}$ a big resolution of W . Note that small resolutions need not be projective, in fact in many cases there exists no projective small resolution. However, $\hat{W}$ is a Moishezon manifold and hence we still have a Hodge decomposition and thus can speak about the Hodge numbers $h^{pq}(\hat{W})$, cf. [Ue]. Let S and S' be the sets of critical values of r and r' . We define $S'' = S \cap S'$ and $\tilde{S} = S \cup S' \setminus S''$. Then all fibres over points in $\tilde{S}$ are of type $I_n \times I_0$ or $I_0 \times I_n$. For any $t \in \mathbb{P}^1$ we denote by $b(t)$, resp. $b'(t)$ the number of components of $r^{-1}(t)$, resp. $r'^{-1}(t)$. Then it is well known [Scho1, p. 190] that

$$h^{12}(\tilde{W}) = h^{12}(\hat{W}) = 1 + \text{rank Pic}(W_\eta) - \#S'' + \sum_{t \in \tilde{S}} (b(t)b'(t) - 1).$$

Here $\text{Pic}(W_\eta)$ is the Picard group of the general fibre. Its rank may be computed using the formula

$$\text{rank Pic}(W_\eta) = \text{rank Pic}(Y_\eta) + \text{rank Pic}(Y'_\eta) + d,$$

where $d = 1$ if Y_η and Y'_η are isogenous and 0 otherwise. By [Scho1, p. 190] this can be rewritten as

$$\text{rank Pic}(W_\eta) = d + 18 - \sum_{t \in S} b(t) - \sum_{t' \in S'} b'(t') + \#S + \#S'.$$

We define the *fibre defect*

$$\delta = \delta(r, r') := 1 + \text{rank Pic}(W_\eta) - \#S''$$

which can be rewritten as

$$\delta = d + 19 - \sum_{t \in S} b(t) - \sum_{t' \in S'} b'(t') + \#S + \#S' - \#S''. \quad (3)$$

When we discuss examples in § 3.2, we will restrict consideration to the case where S and S' are rational elliptic fibrations. Since by Noether's theorem the Euler number of these surfaces is 12 it follows that $\sum_{t \in S} b(t) = \sum_{t' \in S'} b'(t') = 12$, and so in this situation (3) becomes

$$\delta = d - 5 + \#S + \#S' - \#S''. \quad (4)$$

Using its definition, the fibre defect can be computed explicitly for all pairs (r, r') of elliptic fibrations. Note that this number is not an invariant of the variety W but depends on its representation as a fibre product (as we shall see in examples). We call δ the *fibre defect* as this measures the size of $H^{12}(\hat{W}) \oplus H^{21}(\hat{W})$ which does not come from fibres of type $I_n \times I_0$.

We shall now consider the fibres over the points in $\tilde{S}$. As we have already remarked these are of type $I_n \times I_0$ or $I_0 \times I_n$, i.e. a cycle of elliptic ruled surfaces whose components are isomorphic to $\mathbb{P}_i^1 \times E_s$ where $i = 1, \dots, b(t)$, resp. $i = 1, \dots, b'(t)$ and $t \in \tilde{S}$. We shall write $\gamma(t)$ for $b(t)$ or $b'(t)$ in order to simplify the notation. Let $\alpha_t, \beta_t \in H_1(E_t, \mathbb{Z})$ be generators with $\alpha_t^2 = \beta_t^2 = 0$ and $\alpha_t \cdot \beta_t = 1$. Then $A_t^i = \mathbb{P}_i^1 \times \alpha_t$ and $B_t^i = \mathbb{P}_i^1 \times \beta_t$ are generators of $H_3(\mathbb{P}_i^1 \times E_t)$. Finally let $\gamma_{i,t} : \mathbb{P}_i^1 \times E_t \rightarrow \hat{W}$, resp. $\gamma_{i,t} : \mathbb{P}_i^1 \times E_t \rightarrow \tilde{W}$ be the inclusion.

Proposition 2.1. There is an exact sequence

$$0 \rightarrow U \rightarrow H^3(\hat{W}, \mathbb{C}) \xrightarrow{\oplus \gamma_{i,t}^*} \bigoplus_{t \in \tilde{S}} \bigoplus_{i=1}^{\gamma(t)-1} H^3(\mathbb{P}_i^1 \times E_t, \mathbb{C}) \rightarrow 0$$

where $\dim_{\mathbb{C}} U = 2\delta + 2$. A similar exact sequence exists for $\tilde{W}$ instead of $\hat{W}$.

Proof. We have to compute the rank of the intersection matrix of the cycles A_t^i and B_t^i . Clearly cycles in different fibres are independent, hence it suffices to look at one fibre of type $I_n \times I_0$. We assume that the enumeration of the components is such that $\mathbb{P}_i^1 \times E_t$ intersects $\mathbb{P}_j^1 \times E_t$ if and only if $|i - j| \leq 1$ where the index i has to be read cyclically, i.e. we can assume $i \in \mathbb{Z}/\gamma(t)\mathbb{Z}$. Then it follows immediately from the geometry that

$$A_t^i \cdot A_t^j = B_t^i \cdot B_t^j = 0 \text{ for all } i, j$$

and

$$A_t^i . B_t^j = -B_t^j . A_t^i = \begin{cases} 0 & \text{if } |i - j| \geq 2 \\ 1 & \text{if } |i - j| = 1. \end{cases}$$

It remains to determine $A_t^i . B_t^i$. Recall that the normal bundle of a component $\mathbb{P}_i^1 \times E_t$ in $\tilde{W}$, resp. $\tilde{W}$ has degree -2 on the fibres $\mathbb{P}_i^1 \times \{pt\}$. It then follows from [F, 19.2.2] that

$$A_t^i . B_t^i = -B_t^i . A_t^i = -2.$$

Altogether the intersection matrix of the cycles $A_t^i, B_t^i, i = 1, \dots, \gamma(t) - 1$ is given by

$$\begin{pmatrix} \begin{pmatrix} 0 & -2 \\ 2 & 0 \end{pmatrix} & \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} & & & \\ & \begin{pmatrix} 0 & -2 \\ 2 & 0 \end{pmatrix} & \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} & & \\ & & \ddots & \ddots & \\ & & & \begin{pmatrix} 0 & -2 \\ 2 & 0 \end{pmatrix} & \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \\ & & & & \begin{pmatrix} 0 & -2 \\ 2 & 0 \end{pmatrix} \end{pmatrix}.$$

It is obvious that this matrix has rank $2\gamma(t) - 2$ and this proves the proposition. $\square$

Remark 2.2. It is actually not necessary that we start with semi-stable fibrations provided $\tilde{W}$ is a Calabi-Yau variety and we obtain sufficiently many contributions from the fibres of type $I_n \times I_0$.

We now want to consider the modularity of such fibre products.

Theorem 2.3. *Assume that $r : Y \rightarrow \mathbb{P}^1$ and $r' : Y' \rightarrow \mathbb{P}^1$ are semi-stable, rational elliptic fibrations with a section, which are defined over $\mathbb{Q}$. Let $\tilde{W}$ be a (big) resolution of $W = Y \times_{\mathbb{P}^1} Y'$. Moreover assume that*

- (1) *the fibre defect $\delta = \delta(r, r') = 0$,*
- (2) *if $t \in \tilde{S}$ and the fibre W_t is isomorphic to $E_t \times I_n$ for some elliptic curve E_t and an n -cycle $I_n, n \geq 2$, then all components of this fibre are defined over $\mathbb{Q}$.*

Then there is an exact sequence of $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ -modules

$$0 \rightarrow U \rightarrow H_{\text{ét}}^3(\tilde{W}, \mathbb{Q}_l) \rightarrow \bigoplus_{t \in \tilde{S}} \bigoplus_{i=1}^{\gamma(t)-1} H_{\text{ét}}^3(\mathbb{P}_i^1 \times E_t) \rightarrow 0$$

with 2-dimensional kernel U .

Moreover assume that one of the following assumptions holds:

- (3) $\tilde{W}$ has good reduction at 3 and 7 or
- (4) $\tilde{W}$ has good reduction at 5 and some prime $p \equiv \pm 2 \pmod{5}$ with the trace of Frob_p on the kernel U not divisible by 5 or
- (5) $\tilde{W}$ has good reduction at 3 and the trace of Frob_3 on U not divisible by 3.

Then $\tilde{W}$ is modular and the same is true for a small projective resolution defined over $\mathbb{Q}$ if it exists. Moreover

$$L(H_{\text{ét}}^3(\tilde{W}), s) \stackrel{\circ}{=} L(f, s) \prod_{t \in \tilde{S}} (g_{E_t}, s - 1)^{\gamma(t) - 1}$$

where f is a modular form of weight 4 and g_{E_t} is the weight 2 form associated to E_t .

Proof. By Proposition 2.1 the condition $\delta = 0$ assures that the components of the fibres $E_t \times \mathbb{P}^1$ over the points in $\tilde{S}$ span $H^{21}(\tilde{W}) \oplus H^{12}(\tilde{W})$ and that the kernel U has, therefore, dimension 2. The result is now an immediate application of Proposition 1.5. $\square$

3 Examples

In this section we want to discuss various examples where the method described above can be applied. We use (4) to find the fibre defect δ in these examples.

3.1 Relation with previous examples

In [HV] we considered the following examples. Let $\mathcal{E}_{(\alpha, \beta, \gamma)}$ be the elliptic fibration which is the resolution of the surface $\mathcal{E}'_{(\alpha, \beta, \gamma)} \subset \mathbb{P}^2 \times \mathbb{P}^1$ given by

$$\mathcal{E}'_{(\alpha, \beta, \gamma)} : (x + y + z)(\alpha xy + \beta yz + \gamma zx)t_0 = t_1xyz, \quad (5)$$

where $(x : y : z) \in \mathbb{P}^2$, $(t_0 : t_1) \in \mathbb{P}^1$. There is a projection $p : \mathcal{E}_{(\alpha, \beta, \gamma)} \rightarrow \mathbb{P}^1$, with fibre $\mathcal{E}_{(\alpha, \beta, \gamma), t} := p^{-1}(1 : t)$ for generic t . We write $\mathcal{E}_\alpha := \mathcal{E}_{(1, 1, \alpha)}$ and $\mathcal{E}_{\alpha, t} := \mathcal{E}_{(1, 1, \alpha), t}$. The singular fibres of these families can be read off from [HV, Section 4, Table 2].

Taking fibre products and resolving singularities one obtains smooth varieties

$$X_{\mathbf{a}} = X_{(\alpha_1 : \alpha_2 : \alpha_3 : \alpha_4 : \alpha_5 : \alpha_6)} \sim_{\text{bir}} \mathcal{E}_{(\alpha_1 : \alpha_2 : \alpha_3)} \times_{\mathbb{P}^1} \mathcal{E}_{(\alpha_4 : \alpha_5 : \alpha_6)}$$

which are resolutions of nodal Calabi-Yau threefolds. The examples discussed in [HV] were $X_{25} = X_{(1:1:1:1:1:25)}$, $X_{(1:1:1:9:9:9)}$ and $X_{(1:1:1:4:4:9)}$ where $h^{12} = 4, 2$ and 1 respectively. In the last case

$$X_{(1:1:4:4:4:16)} \sim_{bir} \mathcal{E}_{(1:1:16)} \times_{\mathbb{P}^1} \mathcal{E}_{(4:4:4)}$$

and we have $\#S = \#S' = 4$ and $\#S'' = 3$ and hence, since these families are not isogeneous, we have $\delta = 0$. In this case $h^{12} = 1$ and it is sufficient to consider the $I_2 \times I_0$ -fibre over the point 16. We can then derive modularity from Theorem 2.3.

In the second example we can either write

$$X_{(1:1:1:9:9:9)} \sim_{bir} \mathcal{E}_{(1:1:1)} \times_{\mathbb{P}^1} \mathcal{E}_{(9:9:9)}$$

or

$$X_{(1:1:1:9:9:9)} \sim_{bir} \mathcal{E}_{(1:1:9)} \times_{\mathbb{P}^1} \mathcal{E}_{(1:9:9)}.$$

In the first case we have again $\#S = \#S' = 4$ and $\#S'' = 3$ and hence $\delta = 0$. We have a fibre of type $I_3 \times I_0$ over the point 1 and since $h^{12} = 2$ this spans all of $H^{21} \oplus H^{12}$ and our method applies. In the second description we have $\#S = \#S' = 5$ and $\#S'' = 4$ and this gives $\delta = 1$. This is also an example which shows that δ is not a birational invariant but depends on the chosen fibration.

Finally in the first example

$$X_{25} \sim_{bir} \mathcal{E}_{(1:1:1)} \times_{\mathbb{P}^1} \mathcal{E}_{(1:1:25)}$$

and in this case $\#S = 4$, $\#S' = 5$, $\#S'' = 3$ and thus $\delta = 1$. Hence we cannot invoke Theorem 2.3. However, we showed in [HV] that one can still find sufficiently many elliptic ruled surfaces in X_{25} in order to be able to apply Proposition 1.5.

The weight 4 forms corresponding to $H^{30} \oplus H^{03}$ were computed in [HV].

A further class of examples where this method works was discussed by Schütt [Schu2]. In his examples $\#S = 5$ or 4 and $\#S' = 4$ with $\#S'' = 4$ and 3 respectively. In all cases this gives $\delta = 0$ and hence one is in a situation where Theorem 2.3 applies.

3.2 New examples

In this section we give a large number of examples of (birational) Calabi-Yau threefolds for which Theorem 2.3 implies modularity. In many cases we shall also conjecture the modular forms which appear. We have strong numerical evidence for these conjectures, i.e. we computed some of the Fourier coefficients modulo p for all primes of good reduction less than 100. It seems very likely that, using the technique of Faltings, Livné and Serre, these conjectures can be verified.

We use the following notation. If $\mathcal{F}^t$ is an elliptic fibration with parameter t , and if M is a fractional linear transformation, then $\mathcal{F}^{Mt}$ denotes the family obtained by replacing t with Mt .

We denote by $\mathcal{E}(\Gamma)$ Beauville's modular elliptic surfaces [B], where Γ is one of $\Gamma(3)$, $\Gamma_1(4) \cap \Gamma(2)$, $\Gamma_1(6)$, $\Gamma_1(5)$, $\Gamma_0(8) \cap \Gamma_1(4)$ or $\Gamma_0(9) \cap \Gamma_1(3)$. Note that $\mathcal{E}(\Gamma_1(6)) = \mathcal{E}_{(1:1:1)}$. The others may be given by equations as in the table below. The values of t for which the fibres are singular are tabulated, and the last column shows the number of components of the singular fibres, all of which are of I_n type.

Γ	equation for $\mathcal{E}(\Gamma)$	singular t	fibres
$\Gamma(3)$	$x^3 + y^3 + z^3 = 3txyz$	$\infty, 1, \frac{-1 \pm \sqrt{-3}}{2}$	3, 3, 3, 3
$\Gamma_1(4) \cap \Gamma(2)$	$x(x^2 + z^2 + 2zy) = tz(x^2 - y^2)$	$\infty, 0, 1, -1$	4, 4, 2, 2
$\Gamma_1(5)$	$x(x - z)(y - z) = tzy(x - y)$	$\infty, 0, \frac{-11 \pm 5\sqrt{5}}{2}$	5, 5, 1, 1
$\Gamma_1(6)$	$(x + y + z)(xy + yz + zx) = txyz$	$\infty, 0, 1, 9$	6, 2, 3, 1
$\Gamma_0(8) \cap \Gamma_1(4)$	$(x + y)(xy + z^2) = 4txyz$	$\infty, 0, 1, -1$	8, 2, 1, 1
$\Gamma_0(9) \cap \Gamma_1(3)$	$x^2y + y^2z + z^2x = 3txyz$	$\infty, 1, \frac{-1 \pm \sqrt{-3}}{2}$	9, 1, 1, 1

If Y and Y' are **not** isogenous then the condition $\delta = 0$ in Theorem 2.3 translates into

$$\#S + \#S' = \#S'' + 5.$$

This shows that we cannot have examples where at least 1 elliptic fibration has 6 singular fibres. In order to produce examples with $\delta = 0$ we have three possibilities:

	$\#S$	$\#S'$	$\#S''$	possible examples
Case A	5	5	5	$\mathcal{E}_{(1:\alpha^2:\alpha^2)} \times_{\mathbb{P}^1} \mathcal{E}_{(1:\beta^2:\beta^2)}^{Mt}$
Case B	5	4	4	$\mathcal{E}_{(1:\alpha^2:\alpha^2)} \times_{\mathbb{P}^1} \mathcal{E}_{(1:1:1)}^{Mt}$ $\mathcal{E}_{(1:\alpha^2:\alpha^2)} \times_{\mathbb{P}^1} \mathcal{E}^{Mt}(\Gamma)$
Case C	4	4	3	$\mathcal{E}_{(1:1:1)} \times_{\mathbb{P}^1} \mathcal{E}_{(1:1:1)}^{Mt}$ $\mathcal{E}(\Gamma_1) \times_{\mathbb{P}^1} \mathcal{E}^{Mt}(\Gamma_2)$

We now construct explicit examples for each case.

Case A: $\#S = \#S' = \#S'' = 5$ (and the two surfaces are not isogeneous). Notice that this can never give non-rigid examples since we have no fibres of type $I_n \times I_0$.

Given a fibration $\mathcal{E}_{(\alpha,\beta,\gamma)}$ and a fractional linear transformation M , we consider

$$\mathcal{E}_{(\alpha,\beta,\gamma)}^{Mt} : (x + y + z)(axy + byz + czx) = (Mt)xyz.$$

We shall take two families given by $\mathcal{E}_{(1,\alpha^2,\alpha^2)}$ and $\mathcal{E}_{(1,\gamma^2,\gamma^2)}$, with $1, \alpha, 2\alpha, 0$ distinct, and $1, \gamma, 2\gamma, 0$ distinct, and analyse possible fibre products. Recall

			fibre types of $\mathcal{E}_{(1,\gamma^2,\gamma^2)}^{Mt}$									level
α^2	γ^2	Mt	∞	1	0	$(1 \pm 2\alpha)^2$		$(1 \pm 2\alpha)^2$		$(1 \pm 2\gamma)^2$		
$\frac{-1}{8}$	$\frac{-1}{8}$	$1 - t$	I_6	I_2	I_2	I_1	I_1	$\frac{1 \pm \sqrt{-8}}{2}$		$\frac{1 \pm \sqrt{-8}}{2}$		32
$\frac{1}{2}$	$\frac{1}{2}$	$1/t$	I_2	I_6	I_2	I_1	I_1	$3 \pm 2\sqrt{2}$		$3 \pm 2\sqrt{2}$		64
$\frac{1}{9}$	4	$9t$	I_6	I_2	I_1	I_2	I_1	1/9	25/9	9	25	60
$\frac{9}{4}$	$\frac{9}{4}$	$16/t$	I_2	I_6	I_1	I_2	I_1	16	4	16	4	480
$\frac{1}{64}$	$\frac{1}{64}$	$\frac{25}{16} - t$	I_6	I_1	I_1	I_2	I_2	$\frac{9}{16}$	$\frac{25}{16}$	$\frac{9}{16}$	$\frac{25}{16}$	35

Table 1: Rigid Calabi-Yau threefolds of the form $\mathcal{E}_{(1,\alpha^2,\alpha^2)} \times \mathcal{E}_{(1,\gamma^2,\gamma^2)}^{Mt}$

from [HV, Section 4, Table 2] that the singular fibres of $\mathcal{E}_{(1,\alpha^2,\alpha^2)}$ lie over the points $\{\infty, 0, 1, (1 \pm 2\alpha)^2\}$, with fibre types I_6, I_2, I_2, I_1, I_1 respectively, provided $2\alpha \neq 0, 1, 2$. If there is a fractional linear transformation M with $M\{\infty, 0, 1, (1 \pm 2\alpha)^2\} = \{\infty, 0, 1, (1 \pm 2\gamma)^2\}$, then the resolution of

$$\mathcal{E}_{(1,\alpha^2,\alpha^2)} \times \mathcal{E}_{(1,\gamma^2,\gamma^2)}^{Mt}$$

is a rigid (birational) Calabi-Yau threefold, provided that $\#\{\infty, 0, 1, (1 \pm 2\alpha)^2\} = 5$ and that this is not an isogeneous pairing. There are 60 such choices (this can be reduced by symmetry considerations) for which three of the points of $\infty, 0, 1, (1 \pm 2\gamma)^2$ map to 0, 1 and ∞ . This determines M , and the remaining two points β, δ then map to $M(\beta), M(\delta)$, which are given by rational functions in α and γ . The requirement that

$$M(\beta) = (1 + 2\alpha)^2 \text{ and } M(\delta) = (1 - 2\alpha)^2, \quad (6)$$

implies $\alpha = (M(\beta) - M(\delta))/8$. This expression for α in terms of γ can be substituted into (6) to give polynomials satisfied by γ , and from this we can determine whether α^2 and $\gamma^2 \in \mathbb{Q} \setminus \{0\}$. One must also check that $(1 \pm 2\gamma)^2 \neq 0, 1, \infty$. In this case, we get a new example of a rigid Calabi-Yau threefold.

We list all possible cases in Table 1. The singular fibres of the first family have types I_6, I_2, I_2, I_1, I_1 , in this order, so we only list the fibre types of the second family. Note that $\mathcal{E}_{(1,1/9,1/9)}^t \times \mathcal{E}_{(1,4,4)}^{9t}$ is the same as $\mathcal{E}_{(9,1,1)} \times \mathcal{E}_{(1,4,4)} \sim_{bir} X_{(1:1:1:4:4:9)}$, which we considered in [HV], so this case gives only four new cases.

Computational evidence indicates that the L-series of the middle cohomology of $\mathcal{E}_{(1,-1/8,-1/8)}^t \times \mathcal{E}_{(1,-1/8,-1/8)}^{1-t}$ is a weight 4 level 32 modular form, with q -expansion

$$q - 8q^3 - 10q^5 - 16q^7 + 37q^9 + 40q^{11} + \dots$$

Similarly, computations indicate that $\mathcal{E}_{(1, \frac{1}{2}, \frac{1}{2})}^t \times \mathcal{E}_{(1, \frac{1}{2}, \frac{1}{2})}^{1/t}$ is associated with a weight 4 level 64 modular form, with q -expansion

$$q - 22q^5 - 27q^9 + 18q^{13} + \dots,$$

$\mathcal{E}_{(1, 9/4, 9/4)}^t \times \mathcal{E}_{(1, 9/4, 9/4)}^{16/t}$ is associated with a weight 4 level 480 modular form,

$$q + 3q^3 + 5q^5 + 4q^7 + 9q^9 - 40q^{11} - 90q^{13} + \dots,$$

and $\mathcal{E}_{(1, \frac{1}{64}, \frac{1}{64})}^t \times \mathcal{E}_{(1, \frac{1}{64}, \frac{1}{64})}^{25/16-t}$ is associated with the weight 4 level 35 form

$$q + q^2 - 8q^3 - 7q^4 - 5q^5 - 8q^6 + \dots.$$

Case B: $\#S = 5$, $\#S' = \#S'' = 4$.

i. Fibre products of $\mathcal{E}_{(1:a:a)}$ fibrations. We first consider products of the form

$$\mathcal{E}_{(1:\alpha^2:\alpha^2)} \times_{\mathbb{P}^1} \mathcal{E}_{(1:1:1)}^{Mt}$$

for some fractional linear transformation M , and $\alpha \neq 0, 1, 1/2$. Since we are more interested in non-rigid cases, we will not investigate the rigid possibilities. Recall that the singular fibres of $\mathcal{E}_{(1:1:1)}$ lie over the points $\{\infty, 0, 1, 9\}$ and that the singular fibres are of type I_6, I_2, I_3, I_1 (in this order). For a non-rigid example, we are therefore looking for matrices M with

$$\{\infty, 0, 1\} \not\subset M\{\infty, 0, 1, 9\} \subset \{\infty, 0, 1, (1 \pm 2\alpha)^2\}.$$

Denote the cross ratio by $C(a, b, c, d) = (a - c)(b - d)/(a - d)(b - c)$, and set $j(a, b, c, d) = j(C(a, b, c, d))$, where

$$j(\lambda) = \frac{(\lambda^2 - \lambda + 1)^3}{(\lambda - 1)^2 \lambda^2}.$$

Possible values of α^2 are given by solutions to $73^3 2^{-6} 3^{-4} = j(\infty, 0, 1, 9) = J$, for $J = j(0, 1, (1 \pm 2\alpha)^2)$, $j(\infty, 1, (1 \pm 2\alpha)^2)$ or $j(0, \infty, (1 \pm 2\alpha)^2)$. The first possibility,

$$j\left(\left(\frac{2\alpha + 1}{2\alpha - 1}\right)^2 \left(\frac{\alpha + 1}{\alpha - 1}\right)\right) = 73^3 2^{-6} 3^{-4}$$

has no rational solutions. Up to sign, the second,

$$j\left(\frac{\alpha + 1}{\alpha - 1}\right) = \frac{(\alpha^2 + 3)^3}{4(\alpha^2 - 1)^2} = 73^3 2^{-6} 3^{-4}$$

gives $\alpha = 17, 5/4, 7/9$, and the third,

$$j\left(\left(\frac{2\alpha + 1}{2\alpha - 1}\right)^2\right) = 73^3 2^{-6} 3^{-4}$$

α	Mt	fibre types					$(1 \pm 2\alpha)^2$		level	modular form
		∞	1	0	$(1 \pm 2\alpha)^2$					
17	$\frac{(t-1)}{(t-33^2)}$	I_3	I_2	—	I_6	I_1	33^2	35^2	17	$q - 3q^2 - 8q^3 \dots$
$\frac{5}{4}$	$\frac{45}{4(t-1)}$	I_2	I_6	—	I_1	I_3	$\frac{9}{4}$	$\frac{49}{4}$	90	$q - 2q^2 + 4q^4 - 5q^5 - 4q^7 - 8q^8 + 10q^{10} - 12q^{11} \dots$
$\frac{7}{9}$	$\frac{(81t-25)}{56}$	I_6	I_3	—	I_2	I_1	$\frac{25}{81}$	$\frac{23^2}{81}$	21	$q - 3q^2 - 3q^3 \dots$
$\frac{1}{4}$	$\frac{9}{4t}$	I_2	—	I_6	I_1	I_3	$\frac{1}{4}$	$\frac{9}{4}$	90	$q - 2q^2 + 4q^4 + 5q^5 - 4q^7 - 8q^8 - 10q^{10} + 48q^{11} \dots$

Table 2: Modular Calabi-Yau varieties of the form $\mathcal{E}_{(1:\alpha^2:\alpha^2)} \times_{\mathbb{P}^1} \mathcal{E}_{(1:1:1)}^{Mt}$

α^2	Γ	M	level	modular form
−3	$\Gamma(3)$	$(t+7)/8$	27	$q - 3q^2 + q^4 - 15q^5 \dots$
9	$\Gamma_1(4) \cap \Gamma(2)$	$(25-t)/24$	12	$q + 3q^3 - 18q^5 + 8q^7 \dots$
121/125	$\Gamma_1(5)$	$\frac{125}{88}(1-t)$	55	$q + q^2 - 3q^3 - 7q^4 + \dots$
9	$\Gamma_0(8) \cap \Gamma_1(4)$	$24/(25-t)$	12	$q + 3q^3 - 18q^5 + 8q^7 \dots$
−3	$\Gamma_0(9) \cap \Gamma_1(3)$	$(t+7)/8$	27	$q - 3q^2 + q^4 - 15q^5 \dots$

Table 3: Modular Calabi-Yau threefolds of the form $\mathcal{E}_{(1:\alpha^2:\alpha^2)} \times_{\mathbb{P}^1} \mathcal{E}(\Gamma)^{Mt}$

gives $\alpha = 1, 1/4$ (plus irrational solutions), but $\alpha = 1$ is a degenerate case. We obtain the examples given in Table 2. As before, the first family, $\mathcal{E}_{(1:\alpha^2:\alpha^2)}$, has fibre types I_6, I_2, I_2, I_1, I_1 in this order, so we only list the fibre types for the second family, $\mathcal{E}_{(1:1:1)}^{Mt}$. In all cases $h^{12} = 1$. Note that the two level 90 modular forms in this table are not twists of each other.

ii. $\mathcal{E}_{(1:a:a)}$ with Beauville modular fibrations. The same procedure may be applied to fibrations of the form

$$\mathcal{E}_{(1:\alpha^2:\alpha^2)} \times_{\mathbb{P}^1} \mathcal{E}(\Gamma)^{Mt}.$$

Table 3, which is not exhaustive, gives a few examples, all of which have $h^{12} = 1$. Note that the level 55 example is the same as Schütt's W_2 in [Schu2]. There is a 3 : 1 isogeny between the two cases in Table 3 where level 27 occurs. This isogeny is described by Schoen [Scho2].

Case C: $\#S = \#S' = 4$, $\#S'' = 3$.

i. Fibre products of $\mathcal{E}_{(1:1:1)}$ fibrations. We consider non-rigid possibili-

ties of the form

$$\mathcal{E}_{(1:1:1)} \times_{\mathbb{P}^1} \mathcal{E}_{(1:1:1)}^{Mt}.$$

We must take M such that

$$\#(\{0, 1, \infty, 9\} \cup M\{0, 1, \infty, 9\}) = 3.$$

For example, there are 6 possible ways to map $0, 1, 9$ to $0, 1, 9$, only one of which (the identity) maps ∞ to ∞ . The other 5 cases are given by transformations given by matrices

$$M := \begin{pmatrix} 81 & -81 \\ 17 & -81 \end{pmatrix}, \begin{pmatrix} 81 & -81 \\ 73 & -9 \end{pmatrix}, \begin{pmatrix} 9 & -81 \\ 73 & -81 \end{pmatrix}, \begin{pmatrix} 9 & 0 \\ 10 & -9 \end{pmatrix}, \text{ or } \begin{pmatrix} -1 & 9 \\ 7 & 1 \end{pmatrix}.$$

The resulting fibrations all have $h^{12} = 10$.

For each of the 10 choices of pairings, we have 5 maps for which $S \neq S'$ (there are 6 maps, (the number of permutations of three object), but this includes the identity, or a map such as $t \mapsto 9/t$ which leaves the cross ratio of $0, 1, 9, \infty$ invariant (the cross-ratio is invariant under $a \leftrightarrow b, c \leftrightarrow d$). Explicitly, the maps are given by fractional linear transformations defined by matrices $m_i T^j (RT)^k m_l^{-1}$ where $1 \leq i \leq l \leq 4$, $j = 0, 1$, $k = 0, 1, 2$, and $T = \begin{pmatrix} -1 & 1 \\ 0 & 1 \end{pmatrix}$, $R = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, $m_1 = \begin{pmatrix} 9 & 0 \\ 1 & 8 \end{pmatrix}$, $m_2 = \begin{pmatrix} 9 & 0 \\ 0 & 1 \end{pmatrix}$, $m_3 = \begin{pmatrix} 8 & 1 \\ 0 & 1 \end{pmatrix}$, $m_4 = I$. This gives 50 fibrations, (six of which are given in Table 4), which may not all be new, for example,

$$\mathcal{E}_{(1:1:1)} \times \mathcal{E}_{(1:1:1)}^{9t} \sim_{bir} X_{(1:1:1:9:9:9)},$$

and may not all be distinct, for example, $\mathcal{E}_{(1:1:1)} \times \mathcal{E}_{(1:1:1)}^{Mt}$ is the same variety as $\mathcal{E}_{(1:1:1)} \times \mathcal{E}_{(1:1:1)}^{M^{-1}t}$. The only rigid examples are those with $M\{0, 1, \infty\} = \{0, 1, \infty\}$, (two of which are the same) so we have 4 rigid cases, and at most 45 non-rigid cases. All of the rigid cases (given as the first 4 examples in Table 4) have already been studied in detail by Schütt [Schu1]. Schütt also pointed out to us that the level 90 example in Table 4 is a twist of the level 10 example in this table. Note that this level 90 form is the same as the first level 90 example in Table 2, but for that case we had $h^{12} = 1$. We believe that those examples in Table 4 which have the same modular form are isogeneous.

ii. Beauville case.

We can obtain examples from the $\mathcal{E}(\Gamma)$ in exactly the same way as for $\mathcal{E}_{(1:1:1)} = \mathcal{E}(\Gamma_1(6))$ above. However, in general, unless all the singular t are rational, we have fewer cases. Also in some examples, many of the possible M give rise to isomorphic cases. Table 4 gives a number of examples of modular fibrations of the form

$$\mathcal{E}(\Gamma_1) \times_{\mathbb{P}^1} \mathcal{E}^{Mt}(\Gamma_2),$$

Γ_1	Γ_2	M	h^{12}	level	modular form
$\Gamma_1(6)$	$\Gamma_1(6)$	$t/(t-1)$	0	21	$q - 3q^2 - 3q^3 \dots$
$\Gamma_1(6)$	$\Gamma_1(6)$	$1/t$	0	10	$q + 2q^2 - 8q^3 \dots$
$\Gamma_1(6)$	$\Gamma_1(6)$	$1-t$	0	17	$q - 3q^2 - 8q^3 \dots$
$\Gamma_1(6)$	$\Gamma_1(6)$	$(t-1)/t$	0	73	$q + 3q^2 - 8q^3 \dots$
$\Gamma_1(6)$	$\Gamma_1(6)$	$9t$	2	90	$q - 2q^2 + 4q^4 - 5q^5 - 4q^7 - 8q^8 \dots$
$\Gamma_1(6)$	$\Gamma_1(6)$	$9-t$	4	21	$q - 3q^2 - 3q^3 \dots$
$\Gamma_0(8) \cap \Gamma_1(4)$	$\Gamma_0(8) \cap \Gamma_1(4)$	$1-t$	0	12	$q + 3q^3 - 18q^5 \dots$
$\Gamma_1(4) \cap \Gamma(2)$	$\Gamma_1(4) \cap \Gamma(2)$	$1-t$	2	12	$q + 3q^3 - 18q^5 \dots$
$\Gamma_0(8) \cap \Gamma_1(4)$	$\Gamma_0(8) \cap \Gamma_1(4)$	$\frac{-2}{t+1}$	1	48	$q - 3q^3 - 18q^5 \dots$
$\Gamma_0(8) \cap \Gamma_1(4)$	$\Gamma_1(6)$	t	0	80	$q - 2q^3 - 5q^5 \dots$
$\Gamma_1(4) \cap \Gamma(2)$	$\Gamma_1(6)$	t	1	80	$q - 2q^3 - 5q^5 \dots$
$\Gamma_1(4) \cap \Gamma(2)$	$\Gamma_1(6)$	$1/t$	1	80	$q - 2q^3 - 5q^5 \dots$
$\Gamma_1(4) \cap \Gamma(2)$	$\Gamma_1(6)$	$1-t$	1	28	$q - 10q^3 - 8q^5 \dots$
$\Gamma_1(4) \cap \Gamma(2)$	$\Gamma_1(6)$	$1/(1-t)$	1	28	$q - 10q^3 - 8q^5 \dots$
$\Gamma_1(4) \cap \Gamma(2)$	$\Gamma_1(6)$	$1-1/t$	1	68	$q - 2q^3 - 8q^5 \dots$
$\Gamma_0(9) \cap \Gamma_1(3)$	$\Gamma_0(9) \cap \Gamma_1(3)$	$1/t$	16	27	$q - 3q^2 + q^4 \dots$

Table 4: Modular Calabi-Yau threefolds of the form $\mathcal{E}(\Gamma_1) \times \mathcal{E}(\Gamma_2)^{Mt}$

but is far from exhaustive. The level means the conjectural level of the corresponding weight 4 modular form, indicated by computations. Note that there are some isogenous pairs in this table, for which we expect that the weight 4 modular forms are the same. However, h^{12} is not preserved by the map corresponding to the isogeny. Several cases involving $\Gamma_1(4) \cap \Gamma(2)$, with $h^{12} = 1$, could be replaced by rigid cases with $\Gamma_0(8) \cap \Gamma_1(4)$ replacing $\Gamma_1(4) \cap \Gamma(2)$.

The isogenous case: Finally, If Y and Y' are isogenous then $\#S = \#S' = \#S''$, and the condition $\delta = 0$ in Theorem 2.3 translates into $\#S = 4$.

All examples of this type are rigid. This case includes all self fibre products of Beauville examples, given in the appendix to [Y]. Another example is

$$\mathcal{E}(\Gamma_0(8) \cap \Gamma_1(4)) \times_{\mathbb{P}^1} \mathcal{E}^{(t-1)/(t+1)}(\Gamma_0(8) \cap \Gamma_1(4))$$

The fact that $t \mapsto (t-1)/(t+1)$ is an isogeny is explained in more detail in Example 3.5 in [V], which deals with the same elliptic pencil, but with parameter t replaced by $\frac{1}{4t}$.

We have computationally verified that the Mellin transform of the L-series of the middle cohomology of this variety is the weight 4 level 16 mod-

ular form,

$$q + 4q^3 - 2q^5 - 24q^7 - 11q^9 + 44q^{11} \dots$$

References

- [B] A. Beauville, Les familles stables de courbes elliptiques sur $\mathbb{P}^1$ admettant quatre fibres singulières. C. R. Acad. Sci. Paris Sèr. I Math. **294**, no. 19, 657–660 (1982).
- [DM] L. Dieulefait, J. Manoharmayum, Modularity of rigid Calabi-Yau threefolds over $\mathbb{Q}$. In: Calabi-Yau varieties and mirror symmetry (Toronto, ON, 2001), 159–166, Fields Inst. Commun., **38**, Amer. Math. Soc., Providence, RI, 2003.
- [D] L. Dieulefait, From potential modularity to modularity for integral Galois representations and rigid Calabi-Yau threefolds, math.NT/0409102.
- [FM] J. M. Fontaine and W. Messing, p -adic periods and p -adic étale cohomology. In: “Current Trends in Arithmetical Algebraic Geometry (Arcata, Calif., 1985)” . Contemporary Mathematics **67** (1987), 197–207, American Mathematical Society.
- [F] W. Fulton, Intersection Theory, Second Edition, Springer Verlag, 1998.
- [HV] K. Hulek and H. Verrill, On modularity of rigid and non-rigid Calabi-Yau varieties associated to the root lattice A_4 , math.AG/0304169. To appear: Nagoya Math. Journal.
- [LY] R. Livné and N. Yui, The modularity of certain non-rigid Calabi-Yau threefolds, math.AG/0304497.
- [Scho1] C. Schoen, On fiber products of rational elliptic surfaces with section. Math. Z. **197**, 177–199 (1988).
- [Scho2] C. Schoen, On the computation of the cycle class map, Ann. Sci. École Norm. Sup. (4) **28** (1995), no. 1, 1–50
- [Schu1] M. Schütt, New examples of modular rigid Calabi-Yau varieties, Collectanea Mathematica 55,2 (2004), 219–228. (cf. math.AG/0311106.)
- [Schu2] M. Schütt, On the modularity of three Calabi-Yau threefolds with bad reduction at 11, math.AG/0405450. To appear: Bull. Canad. Math. Soc..

- [Ue] K. Ueno, Classification theory of algebraic varieties and compact complex spaces. Lect. Notes Math. **439**, Springer Verlag 1975.
- [V] H. Verrill, Picard-Fuchs equations of some families of elliptic curves, in *Proceedings on Moonshine and related topics (Montréal, 1999)*, CRM Proc. Lecture Notes, 30, 253–268.
- [Y] N. Yui, *Update on the modularity of Calabi-Yau varieties with appendix by Helena Verrill*, Calabi-Yau Varieties and Mirror Symmetry, Fields Institute Communications, Volume: 38 (N. Yui and J. D. Lewis, eds.), 2003.

Klaus Hulek
 Institut für Mathematik (C)
 Universität Hannover
 Welfengarten 1, 30060 Hannover
 Germany
hulek@math.uni-hannover.de

Helena Verrill
 Department of Mathematics
 Louisiana State University
 Baton Rouge, LA 70803-4918
 USA
verrill@math.lsu.edu